

# Skew Hook Formula for $d$ -Complete Posets

Soichi OKADA

(Nagoya University)

joint work with Hiroshi NARUSE (University of Yamanashi)

based on arXiv:1802.09748

Japanese Conference on Combinatorics and its Applications

Sendai, May 21, 2018

## Young Diagrams and Standard Tableaux

For a partition  $\lambda$ , we define its **diagram** by

$$D(\lambda) = \{(i, j) \in \mathbb{Z}^2 : 1 \leq j \leq \lambda_i\}.$$

Let  $\lambda$  and  $\mu$  be partitions such that  $\lambda \supset \mu$  (i.e.,  $D(\lambda) \supset D(\mu)$ ). A **standard tableau** of skew shape  $\lambda/\mu$  is a filling  $T$  of the cells of  $D(\lambda) \setminus D(\mu)$  with numbers  $1, 2, \dots, n = |\lambda| - |\mu|$  satisfying

- each integer appears exactly once,
- the entries in each row and each column are increasing.

### Example

1	2	4	6
3	5	8	
7			

		2	3
1	5	6	
4			

are standard tableaux of shape  $(4, 3, 1)$  and skew shape  $(4, 3, 1)/(2)$  respectively.


## Frame–Robinson–Thrall’s Hook Formulas for Young Diagrams

**Theorem** (Frame–Robinson–Thrall) The number  $f^\lambda$  of standard tableaux of shape  $\lambda$  is given by

$$f^\lambda = \frac{n!}{\prod_{v \in D(\lambda)} h_\lambda(v)}, \quad n = |\lambda|,$$

where  $h_\lambda(i, j) = \lambda_i + \lambda'_j - i - j + 1$  is the **hook length** of  $(i, j)$  in  $D(\lambda)$ .

**Example** The hook of  $(1, 2)$  in  $D(4, 3, 1)$  and the hook lengths are given by



Hence we have

$$f^{(4,3,1)} = \frac{8!}{6 \cdot 4 \cdot 3 \cdot 1 \cdot 4 \cdot 2 \cdot 1 \cdot 1} = 70.$$

## Naruse's Hook Formulas for skew Young Diagrams

**Theorem** (Naruse) The number  $f^{\lambda/\mu}$  of standard tableaux of skew shape  $\lambda/\mu$  is given by


$$f^{\lambda/\mu} = n! \sum_D \frac{1}{\prod_{v \in D(\lambda) \setminus D} h_\lambda(v)}, \quad n = |\lambda| - |\mu|,$$

where  $D$  runs over all excited diagrams of  $D(\mu)$  in  $D(\lambda)$ .

- If a subset  $D \subset D(\lambda)$  and  $u = (i, j)$  satisfy  $(i, j + 1), (i + 1, j), (i + 1, j + 1) \in D(\lambda) \setminus D$ , then we define

$$\alpha_u(D) = D \setminus \{(i, j)\} \cup \{(i + 1, j + 1)\}.$$

- We say that  $D$  is an **excited diagram** of  $D(\mu)$  in  $D(\lambda)$  if  $D$  is obtained from  $D(\mu)$  after a sequence of **elementary excitations**  $D \rightarrow \alpha_u(D)$ .



## Naruse's Hook Formulas for skew Young Diagrams

**Theorem** (Naruse) The number  $f^{\lambda/\mu}$  of standard tableaux of skew shape  $\lambda/\mu$  is given by

$$f^{\lambda/\mu} = n! \sum_D \frac{1}{\prod_{v \in D(\lambda) \setminus D} h_\lambda(v)}, \quad n = |\lambda| - |\mu|,$$

where  $D$  runs over all excited diagrams of  $D(\mu)$  in  $D(\lambda)$ .

**Example** If  $\lambda = (4, 3, 1)$  and  $\mu = (2)$ , then there are three excited diagrams of  $D(\mu)$  in  $D(\lambda)$ :

6	4	3	1
4	2	1	
1			

6	4	3	1
4	2	1	
1			

6	4	3	1
4	2	1	
1			

and we have

$$f^{(4,3,1)/(2)} = 6! \left( \frac{1}{3 \cdot 1 \cdot 4 \cdot 2 \cdot 1 \cdot 1} + \frac{1}{4 \cdot 3 \cdot 1 \cdot 4 \cdot 2 \cdot 1} + \frac{1}{6 \cdot 4 \cdot 3 \cdot 1 \cdot 4 \cdot 1} \right) = 40.$$

## Reverse Plane Partitions

For a poset  $P$ , a  $P$ -partition is a map  $\pi : P \rightarrow \mathbb{N}$  satisfying

$$x \leq y \text{ in } P \implies \pi(x) \geq \pi(y) \text{ in } \mathbb{N}.$$

Let  $\mathcal{A}(P)$  be the set of  $P$ -partitions, and write  $|\pi| = \sum_{x \in P} \pi(x)$  for  $\pi \in \mathcal{A}(P)$ .

The Young diagrams can be regarded as posets by defining

$$(i, j) \geq (i', j') \iff i \leq i', j \leq j'.$$

If  $P = D(\lambda) \setminus D(\mu)$ , then  $P$ -partitions are called **reverse plane partitions** of shape  $\lambda/\mu$ .

### Example

$$\pi = \begin{array}{|c|c|c|c|} \hline & & 3 & 3 \\ \hline 0 & 1 & 3 & \\ \hline 2 & & & \\ \hline \end{array}$$

is a reverse plane partition of shape  $(4, 3, 1)/(2)$  and  $|\pi| = 12$ .

## Univariate Generating Functions of Reverse Plane Partitions

**Theorem** (Stanley) For a partition  $\lambda$ , the generating function of reverse plane partitions of shape  $\lambda$  is given by

$$\sum_{\pi \in \mathcal{A}(D(\lambda))} q^{|\pi|} = \frac{1}{\prod_{v \in P} (1 - q^{h_\lambda(v)})}.$$

**Theorem** (Morales–Pak–Panova) For partitions  $\lambda \supset \mu$ , the generating function of reverse plane partition of skew shape  $\lambda/\mu$  is given by

$$\sum_{\pi \in \mathcal{A}(D(\lambda) \setminus D(\mu))} q^{|\pi|} = \sum_D \frac{\prod_{v \in B(D)} q^{h_\lambda(v)}}{\prod_{v \in D(\lambda) \setminus D} (1 - q^{h_\lambda(v)})},$$

where  $D$  runs over all excited diagrams of  $D(\mu)$  in  $D(\lambda)$ , and  $B(D)$  is the set of **excited peaks** of  $D$ .

Using the theory of  $P$ -partitions, we can derive the hook formula for standard tableaux from those for reverse plane partitions.

## Generalization of Hook Formulas

The Frame–Robinson–Thrall-type hook formula holds for shifted Young diagrams and rooted trees. Proctor introduced a wide class of posets, called  *$d$ -complete posets*.

**Theorem** (Peterson–Proctor) Let  $P$  be a  $d$ -complete poset. Then we can define the *hook lengths*  $h_P(v)$  for  $v \in P$  so that the number of linear extensions of  $P$  is given by

$$\#\{\text{linear extensions of } P\} = \frac{n!}{\prod_{v \in P} h_P(v)}, \quad n = \#P,$$

and the univariate generating function of  $P$ -partitions is given by


$$\sum_{\pi \in \mathcal{A}(P)} q^{|\pi|} = \frac{1}{\prod_{v \in P} (1 - q^{h_P(v)})}.$$

**Goal** Generalize Naruse's and Morales–Pak–Panova's skew hook formulas to  $d$ -complete posets (in other words, generalize Peterson–Proctor's hook formula to skew setting).



## Double-tailed Diamond

- The **double-tailed diamond poset**  $d_k(1)$  ( $k \geq 3$ ) is the poset depicted below:



- A  **$d_k$ -interval** is an interval isomorphic to  $d_k(1)$ .
- A  **$d_k^-$ -convex set** is a convex subset isomorphic to  $d_k(1) - \{\text{top}\}$ .


## $d$ -Complete Posets

**Definition** A finite poset  $P$  is  $d$ -complete if it satisfies the following three conditions for every  $k \geq 3$ :




(D1) If  $I$  is a  $d_k^-$ -convex set, then there exists an element  $v$  such that  $v$  covers the maximal elements of  $I$  and  $I \cup \{v\}$  is a  $d_k$ -interval.

(D2) If  $I = [v, u]$  is a  $d_k$ -interval and  $u$  covers  $w$  in  $P$ , then  $w \in I$ .

(D3) There are no  $d_k^-$ -convex sets which differ only in the minimal elements.



**Example** Shapes (Young diagrams, left), shifted shapes (shifted Young diagrams, middle) and swivels (right) are  $d$ -complete posets.



## Hook Lengths

Let  $P$  be a connected  $d$ -complete poset. For each  $u \in P$ , we define the **hook length**  $h_P(u)$  inductively as follows:


(a) If  $u$  is not the top of any  $d_k$ -interval, then we define

$$h_P(u) = \#\{w \in P : w \leq u\}.$$

(b) If  $u$  is the top of a  $d_k$ -interval  $[v, u]$ , then we define

$$h_P(u) = h_P(x) + h_P(y) - h_P(v),$$

where  $x$  and  $y$  are the sides of  $[v, u]$ .




## Excited Diagrams for $d$ -Complete Posets

Let  $P$  be a connected  $d$ -complete poset.

- We say that  $u \in D$  is  **$D$ -active** if there is a  $d_k$ -interval  $[v, u]$  with  $v \notin D$  such that

$$z \in [v, u] \text{ and } \begin{cases} z \text{ is covered by } u \\ \text{or} \\ z \text{ covers } v \end{cases} \implies z \notin D.$$



- If  $u \in D$  is  $D$ -active, then we define

$$\alpha_u(D) = D \setminus \{u\} \cup \{v\}.$$

Let  $F$  be an order filter of  $P$ .

- We say that  $D$  is an **excited diagram** of  $F$  in  $P$  if  $D$  is obtained from  $F$  after a sequence of **elementary excitations**  $D \rightarrow \alpha_u(D)$ .


## Excited Peaks for $d$ -Complete Posets

Let  $P$  be a  $d$ -complete poset and  $F$  an order filter of  $P$ . To an excited diagram  $D$  of  $F$  in  $P$ , we associate a subset  $B(D) \subset P$ , called the subset of **excited peaks** of  $D$ , as follows:


- (a) If  $D = F$ , then we define  $B(F) = \emptyset$ .
- (b) If  $D' = \alpha_u(D)$  is obtained from  $D$  by an elementary excitation at  $u \in D$ , then

$$B(\alpha_u(D)) = B(D) \setminus \left\{ z \in [v, u] : \begin{array}{l} z \text{ is covered by } u \\ \text{or } z \text{ covers } v \end{array} \right\} \cup \{v\},$$

where  $[v, u]$  is the  $d_k$ -interval with top  $u$ .



**Example** If  $P$  is the Swivel and an order filter  $F$  has two elements, then there are 4 excited diagrams of  $F$  in  $P$ .



Here the shaded cells form an excited diagram and a cell with  $\times$  is an excited peak.

## Main Theorem

**Theorem** (Naruse–Okada) Let  $P$  be a connected  $d$ -complete poset and  $F$  an order filter of  $P$ . Then the univariate generating function of  $(P \setminus F)$ -partitions is given by

$$\sum_{\pi \in \mathcal{A}(P \setminus F)} q^{|\pi|} = \sum_D \frac{\prod_{v \in B(D)} q^{h_P(v)}}{\prod_{v \in P \setminus D} (1 - q^{h_P(v)})},$$

where  $D$  runs over all excited diagrams of  $F$  in  $P$ . More generally, the multivariate generating function of  $(P \setminus F)$ -partitions is given by

$$\sum_{\pi \in \mathcal{A}(P \setminus F)} \prod_{v \in P} \left( z_{c(v)} \right)^{\pi(v)} = \sum_D \frac{\prod_{v \in B(D)} z[H_P(v)]}{\prod_{v \in P \setminus D} (1 - z[H_P(v)])},$$

where  $D$  runs over all excited diagrams of  $F$  in  $P$  and  $z[H_P(v)]$  is the **hook monomial**.



## Main Theorem

**Theorem** (Naruse–Okada) Let  $P$  be a connected  $d$ -complete poset and  $F$  an order filter of  $P$ . Then the multivariate generating function of  $(P \setminus F)$ -partitions is given by

$$\sum_{\pi \in \mathcal{A}(P \setminus F)} \prod_{v \in P} \left( z_{c(v)} \right)^{\pi(v)} = \sum_D \frac{\prod_{v \in B(D)} z[H_P(v)]}{\prod_{v \in P \setminus D} (1 - z[H_P(v)])},$$

where  $D$  runs over all excited diagrams of  $F$  in  $P$ .

### Remark

- If  $F = \emptyset$ , we recover Peterson–Proctor’s hook formula, and our generalization provides an alternate proof.
- If  $P = D(\lambda)$  and  $F = D(\mu)$  are Young diagrams, then the above theorem reduces to Morales–Pak–Panova’s skew hook formula after specializing  $z_i = q$  ( $i \in I$ ).

## Idea of Proof

Given a connected  $d$ -complete poset  $P$ , we can associate the simply-laced Dynkin diagram  $\Gamma$ , the Weyl group  $W$ , the fundamental weight  $\lambda_P$ ,  $\dots$ , and the Kac–Moody partial flag variety  $\mathcal{X}$ . By using the equivariant  $K$ -theory  $K_{\mathcal{T}}(\mathcal{X})$  of  $\mathcal{X}$ , we obtain

$$\xi^v|_w \in \mathbb{Z}[\Lambda] = \bigoplus_{\lambda \in \Lambda} \mathbb{Z}e^\lambda \quad (v, w \in W^{\lambda_P}),$$

where  $\Lambda$  is the weight lattice. Main Theorem follows from

$$\sum_{\pi \in \mathcal{A}(P \setminus F)} z^\pi = \frac{\xi^{w_F}|_{w_P}}{\xi^{w_P}|_{w_P}} = \sum_D \frac{\prod_{v \in B(D)} z[H_P(v)]}{\prod_{v \in P \setminus D} (1 - z[H_P(v)])},$$

where  $z_i = e^{\alpha_i}$  ( $i \in I$ ) and  $w_P$  (resp,  $w_F$ ) is the Weyl group element corresponding to  $P$  (resp.  $F$ ).